Content Warning

Please Note:

This slide deck contains graphic clinical/surgical images of facial and oral injuries. Viewer discretion is advised. Proceed at your own comfort level.

Some slides have been removed in this slide deck from the live presentation.

Oral & Facial Trauma Management

TREASURE VALLEY ORAL & FACIAL SURGERY

Cole W Anderson DMD, MS



None of the planners or presenters for this educational activity have relevant financial relationship(s) to disclose with ineligible companies whose primary business is producing, marketing, selling, re-selling, or distributing healthcare products used by or on patients.

Educational Objectives

- Oral & Facial Trauma Management
- Historic perspective on oral and facial trauma care
- Regional history of facial trauma care
- Accessing emergent services within the Trauma system
- Comprehensive Facial Trauma Management
- Management of Dento-alveolar injuries
- Soft tissue management of oral injuries

An Enigmatic Specialty

- USA: a dental specialty
- Internationally: a medical specialty
- BAOMS, EA Cranio-Maxillofacial Surgery
 - Practicing the full scope of the specialty requires training in both medicine and dentistry




Multi-disciplinary

- OMFS
- ENT
- Plastics

Historical Perspective Facial Trauma

- After lying in a tomb in Thebes, Egypt, for over 3,000 years, the papyrus was sold to the antiquities dealer Edwin Smith in 1862.
- 1906 was donated to 'The New York Historical Society' by his daughter. It was not until...
- 1920 Egyptologist James Henry Breasted for translation.
- The papyrus includes 48 cases of wounds, injuries and fractures



The Local Contract of the cont

\$1 19 ah13. (F.9.)

Historical Perspective Facial Trauma

- Middle ages through Renaissance, Rudimentary basic splinting and bandaging
- Europeans 15th-19th Centuries experimentation on fracture fixation and primitive reconstruction
- WWI was a turning point due to significant progress in many areas of surgery as well as the large scale injuries as a result of modern warfare



Historical Perspective Facial Trauma

- OMFS is a relatively young specialty 1918 AAOMS
- Roots in Dentistry and Medicine
- WWI; Trench warfare, Turning Point
 - 15% with Facial Injuries, unprecedneted numbers
 - Plastic and Jaw Unit 83rd G.H. Wimereux, France
 - Dr. Auguste Charles Valadier, Dental Surgeon
 - British War Office: Commissioned a center for maxillofacial injuries
 - Dr. Harold Gillies (ENT, OMFS, Anest), Dr. Morestin, Dr.
 Ganzer

Facial Surgery Operating Theater



Queens Hospital For Facial Surgery








Gradier K. Mar. Melling Language of hims instruct & Marie Meeting and in the conference of himse Meeting and instruction of the conference of t







USA

- WWI
- Vilray Papin Blair-General Surgeon
 - Organized the Military Facial Trauma Unit
 - Patterned after Dr. Gillies
 - Blair and Robert Ivy OMFS
 - Multidisciplinary Center (ENT, Plast, OMFS, GS, Neuro
 - Transport Prone, Early Tracheostomy, Jaw stabilization

Saint Alphonsus circa 1900




Regional history of facial trauma care

- Saint Alphonsus was the first in Boise est. 1894 by the Sisters of Holy Cross: 5 sisters, 2 doctors and 9 employees
- . St. Luke's 1902, initially 6 bed facility now the largest healt care organization in the state.

Regional history of facial trauma care

- Traditional Hospital Dynamics between Privileging and Obligation. Turf Wars
- Major changes in sub-speciality delivery of care
- Status in 2012-present

Time Sensitve Emergency System



A new perspective

- Altruism and professional ethics
- Watershed case
- Community need
- A compelling priority from the hospital system

Facial Trauma Associates

- . Cole Anderson
- Jeremy Hixsom
- Arman Haghighi
- . Brian Rowan
- . Tim Obrien












Anatomy: Skeletal









Anatomy: Vascular




Anatomy



Anatomy






Patient Evaluation

- Primary Survey
 - Airway
 - Circulation
- Secondary Survery
 - HEENT




Primary Survey

- Airway
 - C-spine immobilization
 - Secure airway
 - Chin Lift
 - Jaw thrust
 - OPA
 - Nasal airway



Airway Management Oropharyngeal Airway





Airway Management Nasopharyngeal Airway



Definitive Airway

- Intubation
 - Oral
 - Nasal
 - Submental











Primary Survey

- Surgical Airway
 - Cricothyroidotomy
 - Tracheostomy



Primary Survey

- Circulation
 - Restoration of blood volume + C.O.
 - Control of Bleeding
 - Vascular injury
 - Epistaxis




Vascular Injuries

- Penetrating head and neck trauma
- Angiography
- Plain films
- CT
- MRI





Anterior & Posterior Nasal Packs




Neck Injuries

Zone III

Zone II


Zone I




Secondary Survey

- History
 - Mechanism of injury
 - High Energy Injuries
 - MVA
 - Gun Shot Wounds







- History
 - Mechanism of injury
 - Low-Moderate Energy Injuries
 - Assaults
 - Sports Related













CT Images

- Low Energy
 - Fall from standing height
 - Fist to face
- Moderate Energy
 - Crash; bicycle, skate board
 - Baseball bat, hammer
- High Energy
 - MVC
 - GSW



Low energy

- Soft tissue injury; contusion, laceration
- Hard tissue injury; minimal comminution
 - Minimally to non-displaced fractures
- Limited concominant injuries





Moderate Energy

- Soft tissue; can be avulsive, generalized abrasions
- Hard tissue; increased comminutia, vector of energy observed with displaced bone segments
- Concominant injuries; Intra-cranial injury, ocular injury

High Energy

- Soft Tissue: avulsive, crush injuries, lacerations
- Collapse of the buttress of the facial skeleton
- Facial width increase
- High degree of comminutia, Pan-facial fractures
- Skull base and intra-cranial injury




















- Physical Examination
 - Periorbital Ecchymosis
 - Battle's sign
 - Subconjunctival Heme
 - Facial Deformity
 - Occlusal step
 - Numbness








- Pitfalls:
 - Pending airway obstruction
 - Changes in airway status
 - Expanding hematoma
 - Loss of mandibular support of airway
 - Exsanguinating fractures
 - Lacrimal duct injuries
 - Facial nerve injuries

- Cranial Nerve Examination
 - CN I Olfactory Nerve
 - CN II Optic Nerve
 - CN III Oculomotor Nerve
 - CN IV Trochlear Nerve
 - CN V Trigeminal Nerve
 - CN VII Facial Nerve


CN I – Olfactory Nerve

- Sense of Smell
- Cribiform Plate
- Injured with LeFort II and III Fractures




CN II – Optic Nerve

- Function Vision
- Sensory limb of Pupillary Reflex
- Dysfunction may be associated with Orbital or Midface Fractures



CN III – Oculomotor Nerve


- Extraocular Muscles
 - Medial Rectus
 - Superior Rectus
 - Inferior Rectus
 - Inferior Oblique muscle
- Pupillary Constriction
- Eyelid Elevation
- May be injured in orbital or midfacial fractures






CN IV

- CN IV Trochlear
 - Superior Oblique Muscle
 - Injury Inability to gaze inferio-medially




CN V – Trigeminal Nerve

- CN V Trigeminal Nerve
 - Sensory Nerve
 - Injured in blunt trauma associated with facial fractures



CN VI – Abducens Nerve


- CN VI Abducens
 - Lateral Rectus Muscle
 - Long intracranial course
 - Injury inability to gaze laterally





CN VII – Facial Nerve

- CN VII Facial Nerve
 - Motor nerve which moves face
 - Most commonly injured in penetrating Trauma
 - Injury causes unilateral paralysis of facial muscles









В


Principles of Craniofacial Trauma Surgery





Facial Buttresses


- Anterior Buttress (height)
 - Nasomaxillary
 - Zygomaticomaxillary
- Horizontal (Transverse)
 - Alveolus/Palate
 - Pyriform Aperature
 - Orbital Rims
- Posterior Buttress (Height)
 - Pterygomaxillary
- Sagittal Buttress(Projection)
 - Zygomatic Arch



Principles of Craniofacial Trauma Surgery


- Anatomic Reduction
 - Intermaxillary fixation






Principles of Cranio-Maxillofacial Trauma Surgery

- Surgical Exposure
 - Coronal Flap
 - Lefort II, III
 - ZMC
 - Transconj + Subciliary
 - ZMC
 - Orbital Floor Fractures



Principles of Craniofacial Trauma Surgery

- Surgical Exposure
 - Maxillary Vestibular
 - Lefort I
 - ZMC
 - Mandibular Vestibular
 - Mandibular Fractures




Categorized fractures Top to Bottom

- Frontal Sinus
- NOE
- Orbital Fracture
- ZMC
- Lefort Fractures
- Mandibular
 Fractures

Frontal Sinus Fractures







- Usually high energy injury with Periorbital ecchymosis and edema
- CSF leak Sinus precautions, Antibiotics???
- Treatment within 7-10 days

Frontal Sinus Fractures






Frontal Sinus Fractures





Orbital Fractures







- Presentation orbital pain, periorbital ecchymosis and edema, visual changes, enophthalmos, restriction of gaze, palpable rim deformity
- Ophthalmology Consultation???
- Compartment Syndrome; emergent decompression, anticoagulation
- Closely evaluate vision, extraocular muscle function
 - Sinus precautions Antibiotics ???

 - Surgery 7-14 days


Orbital Fractures







Indications for surgery

- Peri-orbital Entrapment
 - Diplopia, restriction of gaze
- Compartment Syndrome
 - Retrobulbar hematoma
 - Anticoagulation
 - Blow out Vs. Trap door
- Expanded orbital volume
 - Enopthalmus risk




Zygomatic Complex Fractures






- Presentation flattened malar eminence, periorbital ecchymosis, paresthesia of cheek, upper lip(CN V), enophthalmos, rim step
- High energy injuries Evaluate globe, vision closely
 - Sinus precaution; Air emphysema
- Treatment 7-10 days








LeFort I









- High energy injury with periorbital ecchymosis, edema, malocclusion
- Transverse Maxillary Fracture detaches tooth bearing segment of maxilla
 - Sinus precautions
- Airway generally intact
- Treatment within 7-10 days
- Change in Occlusion (bite)
- V2 paresthesia, numb teeth

LeFort II Fracture









- High energy injury with periorbital ecchymosis, edema, malocclusion
 - Pyramidal dysjunction floating midface
- May involve cribiform plate, CSF leak, Olfactory Nerve(CN I), Head injury
 - Sinus precautions
- Airway needs to be evaluated closely
- V2 paresthesia, numb teeth
- Treatment within 7-10 days

LeFort III + NOE Fractures







- High energy injury with periorbital ecchymosis, edema, malocclusion

 Craniofacial dysjunction floating midface


 Increased inter-canthal distance

 - Saddle nose deformity
- May involve cribiform plate, CSF leak, Olfactory Nerve(CN I), Head injury Sinus precautions
- Airway needs to be evaluated closely
- Treatment within 7-10 days

Nasal Bone/Septum Fracture

Acute Nasal Fractures


- Management
 - Closed reduction at 5-10 days
 - Anesth. deep MAC or General
 - reduction and elevation





Mandibular Fractures






- Usually present with malocclusion, paresthesia(CN V)
 - Evaluate for hematoma floor of mouth, if flail mandible evaluate airway patency
 - Missing teeth or displaced fracture segments
- High energy injuries evaluate for other fractures
- Treatment within 1-5 days(unless temporarily stabilized)













DENTOALVEOLAR TRAUMA

TRAUMA

» CLASSIFICATION (ANDREASEN)

- CROWN INFRACTION
- UNCOMPLICATED CROWN FRACTURE
- COMPLICATED CROWN FRACTURE
- CROWN AND ROOT FRACTURE
- HORIZONTAL ROOT FRACTURE
- CONCUSSION
- SUBLUXATION
- LATERAL LUXATION
- EXTRUSIVE LUXATION
- INTRUSIVE LUXATION
- AVULSION

TRAUMA

» CLASSIFICATION (ELLIS)

- Type I: Enamel only
- Type II: Enamel and dentin
- Type III: Pulp
- Type IV: Root fracture
- Type V: Luxation
- Type VI: Avulsion

HORIZONTAL ROOT FRACTURE







CONCUSSION

- » Tooth is tender to percussion but no mobility.
- » Baseline pulp test (3% nonvital).
- » Look for root fracture.
- » Soft diet, rinse with chlorhexidine every 12 hrs for 1 week.
- » No treatment required, monitor for 2-3 wks and then regular follow-ups.



SUBLUXATION

- » Some mobility but no displacement.
- » Baseline pulp test (6% nonvital).
- » Look for root fracture.
- » Soft diet, rinse with chlorhexidine every 12 hrs for 1 week.
- » No treatment required, monitor for 2-3 wks and then regular follow-ups.



EXTRUSIVE AND LATERAL LUXATION





LATERAL AND EXTRUSIVE LUXATION

- » Tooth is displaced laterally or coronally.
- » Baseline pulp test:
 - Lateral Luxation: 58% nonvital.
 - Extrusive Luxation: 26% nonvital.

» Treatment:

- Flexible splint up to 3 weeks.
- Soft diet, rinse with chlorhexidine every 12 hrs for 1 week.
- If no response at 2-3 weeks perform RCT.
- Follow-up at 3, 6, 12 months.

INTRUSIVE LUXATION

- » Displacement towards the socket.
- » Baseline pulp test (85% nonvital).
- » Treatment:
 - Soft diet, rinse with chlorhexidine every 12 hrs for 1 week.
 - Immature root formation: May reerupt on its own, if not surgical or orthodontic movement. If no response after 3 wks perform RCT.
 - Mature root formation: Surgical or orthodontic movement, followed by RCT.



- » Tooth is displaced from the socket.
- » Treatment is based on:
 - Stage of root formation
 - Dry time
- » Transport medium:
 - HBSS
 - Skim Milk
- » Antibiotics:
 - Pen VK
 - Doxycycline



- » TREATMENT (Trope and Andreasen)
 - Dry Time less than 1 hour (OPEN APICES)
 - Revascularization of the pulp is good.
 - Place tooth in transport medium.
 - Clean foramen with saline and soak tooth in 100mg doxycycline/20mg saline.
 - Flexible splint 7 to 10 days.
 - Soft diet, rinse with chlorhexidine every 12 hrs for 1 week.
 - Monitor pulp status, may take up to 3 months for the pulp to respond to cold.

» TREATMENT

- Dry Time of less than 1 hour (CLOSED APICES)
 - Place tooth in transport medium.
 - No chance for revascularization.
 - Rinse socket and remove coagulum and gently replace tooth.
 - Flexible splint for 7 to 10 days.
 - Soft diet, rinse with chlorhexidine every 12 hrs for 1 week.
 - RCT in 7 to 10 days with CaOH to decrease resorption.

» TREATMENT

- Dry Time of greater than 1 hour (OPEN APICES)
 - No chance of revascularization.
 - Resorption will occur.
 - Replantation is not recommended.

» TREATMENT

- Dry Time of greater than 1 hour (CLOSED APICES)
 - All PDL cells are dead, no reason to soak in HBSS or Viaspan.
 - To decrease resorption soak tooth in available fluoride solution for at least 5 min.
 - Flexible splint for 7 to 10 days.
 - Soft diet, rinse with chlorhexidine every 12 hrs for 1 week.
 - RCT in 7 to 10 days with CaOH to decrease resorption.
 - Prognosis: guarded

ALVEOLAR FRACTURES

» Alveolar fractures:

- Entire segment moves.
- X-ray.
- Reposition segment and place rigid or semi-rigid splint for 4 weeks.
- Placing a rigid splint increases the chance of resorption and should only be placed when an alveolar fracture is present.




Alveolar Process Injuries

Fractured teeth

Fractured roots

Avulsed teeth

Alveolar fractures
Intruded
Extruded
Displaced












Composite bonding an arch wire to stabilize avulsed teeth and / or alveolar fractures
























Dento-alveolar Case

